

## Useful data

$A \setminus B$	$\{a \in A : a \notin B\}$
$\mathbb{C}$	Set of all complex numbers
$\mathbb{C}^{m \times n}$	Set of all matrices of order $m \times n$ with complex entries
$\mathbb{C}^\infty(\Omega)$	Collection of all infinitely differentiable functions on the open domain $\Omega$
$i$	$\sqrt{-1}$
$I$	Identity matrix of appropriate order
$L^2(\mathbb{R})$	$:= L^2(\mathbb{R}, dx)$
$L^2[a, b]$	$:= L^2([a, b], dx)$
$\mathbb{N}$	Set of all positive integers
$\mathbb{Q}$	Set of all rational numbers
$\mathbb{R}$	Set of all real numbers
$\mathbb{R}^{m \times n}$	Set of all matrices of order $m \times n$ with real entries
$\mathbb{S}^1$	$\{(x_1, x_2) \in \mathbb{R}^2 : x_1^2 + x_2^2 = 1\}$
$\mathbb{S}^2$	$\{(x_1, x_2, x_3) \in \mathbb{R}^3 : x_1^2 + x_2^2 + x_3^2 = 1\}$
$\mathbb{Z}$	Set of all integers

## GATE 2022 General Aptitude (GA)

**Q.1 – Q.5 Carry ONE mark each.**

Q.1	As you grow older, an injury to your _____ may take longer to _____.
(A)	heel / heel
(B)	heal / heel
(C)	heal / heal
(D)	heel / heal


---

Q.2	<p>In a 500 m race, P and Q have speeds in the ratio of 3 : 4. Q starts the race when P has already covered 140 m.</p> <p>What is the distance between P and Q (in m) when P wins the race?</p>
(A)	20
(B)	40
(C)	60
(D)	140

Q.3	<p>Three bells P, Q, and R are rung periodically in a school. P is rung every 20 minutes; Q is rung every 30 minutes and R is rung every 50 minutes.</p> <p>If all the three bells are rung at 12:00 PM, when will the three bells ring together again the next time?</p>
(A)	5:00 PM
(B)	5:30 PM
(C)	6:00 PM
(D)	6:30 PM

---

Q.4	<p>Given below are two statements and four conclusions drawn based on the statements.</p> <p>Statement 1: Some bottles are cups.</p> <p>Statement 2: All cups are knives.</p> <p>Conclusion I: Some bottles are knives.</p> <p>Conclusion II: Some knives are cups.</p> <p>Conclusion III: All cups are bottles.</p> <p>Conclusion IV: All knives are cups.</p> <p>Which one of the following options can be logically inferred?</p>
(A)	Only conclusion I and conclusion II are correct
(B)	Only conclusion II and conclusion III are correct
(C)	Only conclusion II and conclusion IV are correct
(D)	Only conclusion III and conclusion IV are correct

<p>Q.5</p>	<p>The figure below shows the front and rear view of a disc, which is shaded with identical patterns. The disc is flipped once with respect to any one of the fixed axes 1-1, 2-2 or 3-3 chosen uniformly at random.</p> <p>What is the probability that the disc <b>DOES NOT</b> retain the same front and rear views after the flipping operation?</p> <div style="text-align: center;">  <div style="display: flex; justify-content: space-around; width: 100%;"> <span>Front View</span> <span>Rear View</span> </div> </div>
(A)	0
(B)	$\frac{1}{3}$
(C)	$\frac{2}{3}$
(D)	1

**Q. 6 – Q. 10 Carry TWO marks each.**

Q.6	<p>Altruism is the human concern for the wellbeing of others. Altruism has been shown to be motivated more by social bonding, familiarity and identification of belongingness to a group. The notion that altruism may be attributed to empathy or guilt has now been rejected.</p> <p>Which one of the following is the CORRECT logical inference based on the information in the above passage?</p>
(A)	Humans engage in altruism due to guilt but not empathy
(B)	Humans engage in altruism due to empathy but not guilt
(C)	Humans engage in altruism due to group identification but not empathy
(D)	Humans engage in altruism due to empathy but not familiarity

Q.7	<p>There are two identical dice with a single letter on each of the faces. The following six letters: Q, R, S, T, U, and V, one on each of the faces. Any of the six outcomes are equally likely.</p> <p>The two dice are thrown once independently at random.</p> <p>What is the probability that the outcomes on the dice were composed only of any combination of the following possible outcomes: Q, U and V?</p>
(A)	$\frac{1}{4}$
(B)	$\frac{3}{4}$
(C)	$\frac{1}{6}$
(D)	$\frac{5}{36}$



Q.8	<p>The price of an item is 10% cheaper in an online store S compared to the price at another online store M. Store S charges ₹ 150 for delivery. There are no delivery charges for orders from the store M. A person bought the item from the store S and saved ₹ 100.</p> <p>What is the price of the item at the online store S (in ₹) if there are no other charges than what is described above?</p>
(A)	2500
(B)	2250
(C)	1750
(D)	1500

Q.9	<p>The letters P, Q, R, S, T and U are to be placed one per vertex on a regular convex hexagon, but not necessarily in the same order.</p> <p>Consider the following statements:</p> <ul style="list-style-type: none"> <li>• The line segment joining R and S is longer than the line segment joining P and Q.</li> <li>• The line segment joining R and S is perpendicular to the line segment joining P and Q.</li> <li>• The line segment joining R and U is parallel to the line segment joining T and Q.</li> </ul> <p>Based on the above statements, which one of the following options is CORRECT?</p>
(A)	The line segment joining R and T is parallel to the line segment joining Q and S
(B)	The line segment joining T and Q is parallel to the line joining P and U
(C)	The line segment joining R and P is perpendicular to the line segment joining U and Q
(D)	The line segment joining Q and S is perpendicular to the line segment joining R and P

Q.10	<div data-bbox="738 205 1031 451" data-label="Image"> <p>A 4x4 grid of squares. The bottom-left square is labeled 'P' at its bottom-left corner.</p> </div> <p>An ant is at the bottom-left corner of a grid (point P) as shown above. It aims to move to the top-right corner of the grid. The ant moves only along the lines marked in the grid such that the current distance to the top-right corner strictly decreases.</p> <p>Which one of the following is a part of a possible trajectory of the ant during the movement?</p>
(A)	<div data-bbox="406 898 527 1003" data-label="Image"> <p>A path starting at point P, moving right, then up, then right, then up, then right.</p> </div>
(B)	<div data-bbox="414 1102 544 1207" data-label="Image"> <p>A path starting at point P, moving up, then right, then up, then right, then up.</p> </div>
(C)	<div data-bbox="406 1306 527 1411" data-label="Image"> <p>A path starting at point P, moving up, then right, then up, then right.</p> </div>
(D)	<div data-bbox="406 1509 479 1572" data-label="Image"> <p>A path starting at point P, moving right, then up.</p> </div>

**GATE 2022 Mathematics (MA)**

**Q.11 – Q.35 Carry ONE mark each.**

Q.11	<p>Suppose that the characteristic equation of <math>M \in \mathbb{C}^{3 \times 3}</math> is</p> $\lambda^3 + \alpha\lambda^2 + \beta\lambda - 1 = 0,$ <p>where <math>\alpha, \beta \in \mathbb{C}</math> with <math>\alpha + \beta \neq 0</math>. Which of the following statements is TRUE?</p>
(A)	$M(I - \beta M) = M^{-1}(M + \alpha I)$
(B)	$M(I + \beta M) = M^{-1}(M - \alpha I)$
(C)	$M^{-1}(M^{-1} + \beta I) = M - \alpha I$
(D)	$M^{-1}(M^{-1} - \beta I) = M + \alpha I$

# GATE 2022 Mathematics (MA)

Q.12	<p>Consider</p> <p><b>P:</b> Let <math>M \in \mathbb{R}^{m \times n}</math> with <math>m &gt; n \geq 2</math>. If <math>\text{rank}(M) = n</math>, then the system of linear equations <math>Mx = 0</math> has <math>x = 0</math> as the only solution.</p> <p><b>Q:</b> Let <math>E \in \mathbb{R}^{n \times n}, n \geq 2</math> be a non-zero matrix such that <math>E^3 = 0</math>. Then <math>I + E^2</math> is a singular matrix.</p> <p>Which of the following statements is TRUE?</p>
(A)	Both <b>P</b> and <b>Q</b> are TRUE
(B)	Both <b>P</b> and <b>Q</b> are FALSE
(C)	<b>P</b> is TRUE and <b>Q</b> is FALSE
(D)	<b>P</b> is FALSE and <b>Q</b> is TRUE

# GATE 2022 Mathematics (MA)

Q.13	<p>Consider the real function of two real variables given by</p> $u(x, y) = e^{2x}[\sin 3x \cos 2y \cosh 3y - \cos 3x \sin 2y \sinh 3y].$ <p>Let <math>v(x, y)</math> be the harmonic conjugate of <math>u(x, y)</math> such that <math>v(0, 0) = 2</math>. Let <math>z = x + iy</math> and <math>f(z) = u(x, y) + iv(x, y)</math>, then the value of <math>4 + 2if(i\pi)</math> is</p>
(A)	$e^{3\pi} + e^{-3\pi}$
(B)	$e^{3\pi} - e^{-3\pi}$
(C)	$-e^{3\pi} + e^{-3\pi}$
(D)	$-e^{3\pi} - e^{-3\pi}$

**GATE 2022 Mathematics (MA)**

Q.14	<p>The value of the integral</p> $\int_C \frac{z^{100}}{z^{101} + 1} dz$ <p>where <math>C</math> is the circle of radius 2 centred at the origin taken in the anti-clockwise direction is</p>
(A)	$-2\pi i$
(B)	$2\pi$
(C)	0
(D)	$2\pi i$

**GATE 2022 Mathematics (MA)**

Q.15	Let $X$ be a real normed linear space. Let $X_0 = \{x \in X : \ x\  = 1\}$ . If $X_0$ contains two distinct points $x$ and $y$ and the line segment joining them, then, which of the following statements is TRUE?
(A)	$\ x + y\  = \ x\  + \ y\ $ and $x, y$ are linearly independent
(B)	$\ x + y\  = \ x\  + \ y\ $ and $x, y$ are linearly dependent
(C)	$\ x + y\ ^2 = \ x\ ^2 + \ y\ ^2$ and $x, y$ are linearly independent
(D)	$\ x + y\  = 2\ x\ \ y\ $ and $x, y$ are linearly dependent



**GATE 2022 Mathematics (MA)**

Q.16	<p>Let <math>\{e_k : k \in \mathbb{N}\}</math> be an orthonormal basis for a Hilbert space <math>H</math>.</p> <p>Define <math>f_k = e_k + e_{k+1}, k \in \mathbb{N}</math> and <math>g_j = \sum_{n=1}^j (-1)^{n+1} e_n, j \in \mathbb{N}</math>.</p> <p>Then <math>\sum_{k=1}^{\infty}  \langle g_j, f_k \rangle ^2 =</math></p>
(A)	0
(B)	$j^2$
(C)	$4j^2$
(D)	1

**GATE 2022 Mathematics (MA)**

Q.17	Consider $\mathbb{R}^2$ with the usual metric. Let $A = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 \leq 1\}$ and $B = \{(x, y) \in \mathbb{R}^2 : (x-2)^2 + y^2 \leq 1\}$ . Let $M = A \cup B$ and $N = \text{interior}(A) \cup \text{interior}(B)$ . Then, which of the following statements is TRUE?
(A)	$M$ and $N$ are connected
(B)	Neither $M$ nor $N$ is connected
(C)	$M$ is connected and $N$ is not connected
(D)	$M$ is not connected and $N$ is connected

**GATE 2022 Mathematics (MA)**

Q.18	<p>The real sequence generated by the iterative scheme</p> $x_n = \frac{x_{n-1}}{2} + \frac{1}{x_{n-1}}, \quad n \geq 1$
(A)	converges to $\sqrt{2}$ , for all $x_0 \in \mathbb{R} \setminus \{0\}$
(B)	converges to $\sqrt{2}$ , whenever $x_0 > \sqrt{\frac{2}{3}}$
(C)	converges to $\sqrt{2}$ , whenever $x_0 \in (-1, 1) \setminus \{0\}$
(D)	diverges for any $x_0 \neq 0$

**GATE 2022 Mathematics (MA)**

Q.19	<p>The initial value problem</p> $\frac{dy}{dx} = \cos(xy), \quad x \in \mathbb{R}, \quad y(0) = y_0,$ <p>where <math>y_0</math> is a real constant, has</p>
(A)	a unique solution
(B)	exactly two solutions
(C)	infinitely many solutions
(D)	no solution

**GATE 2022 Mathematics (MA)**

Q.20	<p>If eigenfunctions corresponding to distinct eigenvalues <math>\lambda</math> of the Sturm-Liouville problem</p> $\frac{d^2y}{dx^2} - 3\frac{dy}{dx} = \lambda y, \quad 0 < x < \pi,$ $y(0) = y(\pi) = 0$ <p>are orthogonal with respect to the weight function <math>w(x)</math>, then <math>w(x)</math> is</p>
(A)	$e^{-3x}$
(B)	$e^{-2x}$
(C)	$e^{2x}$
(D)	$e^{3x}$

**GATE 2022 Mathematics (MA)**

Q.21	<p>The steady state solution for the heat equation</p> $\frac{\partial u}{\partial t} - \frac{\partial^2 u}{\partial x^2} = 0, \quad 0 < x < 2, \quad t > 0,$ <p>with the initial condition <math>u(x, 0) = 0</math>, <math>0 &lt; x &lt; 2</math> and the boundary conditions <math>u(0, t) = 1</math> and <math>u(2, t) = 3</math>, <math>t &gt; 0</math>, at <math>x = 1</math> is</p>
(A)	1
(B)	2
(C)	3
(D)	4

# GATE 2022 Mathematics (MA)

Q.22	<p>Consider <math>([0, 1], T_1)</math>, where <math>T_1</math> is the subspace topology induced by the Euclidean topology on <math>\mathbb{R}</math>, and let <math>T_2</math> be <i>any</i> topology on <math>[0, 1]</math>. Consider the following statements:</p> <p><b>P</b> : If <math>T_1</math> is a proper subset of <math>T_2</math>, then <math>([0, 1], T_2)</math> is not compact.</p> <p><b>Q</b> : If <math>T_2</math> is a proper subset of <math>T_1</math>, then <math>([0, 1], T_2)</math> is not Hausdorff.</p> <p>Then</p>
(A)	<b>P</b> is TRUE and <b>Q</b> is FALSE
(B)	Both <b>P</b> and <b>Q</b> are TRUE
(C)	Both <b>P</b> and <b>Q</b> are FALSE
(D)	<b>P</b> is FALSE and <b>Q</b> is TRUE

**GATE 2022 Mathematics (MA)**

Q.23	Let $p : ([0, 1], T_1) \rightarrow (\{0, 1\}, T_2)$ be the quotient map, arising from the characteristic function on $[\frac{1}{2}, 1]$ , where $T_1$ is the subspace topology induced by the Euclidean topology on $\mathbb{R}$ . Which of the following statements is TRUE?
(A)	$p$ is an open map but not a closed map
(B)	$p$ is a closed map but not an open map
(C)	$p$ is a closed map as well as an open map
(D)	$p$ is neither an open map nor a closed map



**GATE 2022 Mathematics (MA)**

Q.24	<p>Set <math>X_n := \mathbb{R}</math> for each <math>n \in \mathbb{N}</math>. Define <math>Y := \prod_{n \in \mathbb{N}} X_n</math>. Endow <math>Y</math> with the product topology, where the topology on each <math>X_n</math> is the Euclidean topology. Consider the set</p> $\Delta = \{(x, x, x, \dots) \mid x \in \mathbb{R}\}$ <p>with the subspace topology induced from <math>Y</math>. Which of the following statements is TRUE?</p>
(A)	$\Delta$ is open in $Y$
(B)	$\Delta$ is locally compact
(C)	$\Delta$ is dense in $Y$
(D)	$\Delta$ is disconnected

**GATE 2022 Mathematics (MA)**

Q.25	<p>Consider the linear system of equations <math>Ax = b</math> with</p> $A = \begin{pmatrix} 3 & 1 & 1 \\ 1 & 4 & 1 \\ 2 & 0 & 3 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} 2 \\ 3 \\ 4 \end{pmatrix}.$ <p>Which of the following statements are TRUE?</p>
(A)	The Jacobi iterative matrix is $\begin{pmatrix} 0 & 1/4 & 1/3 \\ 1/3 & 0 & 1/3 \\ 2/3 & 0 & 0 \end{pmatrix}$
(B)	The Jacobi iterative method converges for any initial vector
(C)	The Gauss-Seidel iterative method converges for any initial vector
(D)	The spectral radius of the Jacobi iterative matrix is less than 1

**GATE 2022 Mathematics (MA)**

Q.26	The number of non-isomorphic abelian groups of order $2^2 \cdot 3^3 \cdot 5^4$ is _____.

**GATE 2022 Mathematics (MA)**

Q.27	The number of subgroups of a cyclic group of order 12 is _____.

## GATE 2022 Mathematics (MA)

Q.28	<p>The radius of convergence of the series</p> $\sum_{n \geq 0} 3^{n+1} z^{2n}, \quad z \in \mathbb{C}$ <p>is _____ (round off to TWO decimal places).</p>

**GATE 2022 Mathematics (MA)**

Q.29	<p>The number of zeros of the polynomial</p> $2z^7 - 7z^5 + 2z^3 - z + 1$ <p>in the unit disc <math>\{z \in \mathbb{C} :  z  &lt; 1\}</math> is _____.</p>

**GATE 2022 Mathematics (MA)**

Q.30	<p>If <math>P(x)</math> is a polynomial of degree 5 and</p> $\alpha = \sum_{i=0}^6 P(x_i) \left( \prod_{j=0, j \neq i}^6 (x_i - x_j)^{-1} \right),$ <p>where <math>x_0, x_1, \dots, x_6</math> are distinct points in the interval <math>[2, 3]</math>, then the value of <math>\alpha^2 - \alpha + 1</math> is _____.</p>

**GATE 2022 Mathematics (MA)**

Q.31	The maximum value of $f(x, y) = 49 - x^2 - y^2$ on the line $x + 3y = 10$ is _____.



**GATE 2022 Mathematics (MA)**

Q.32	If the function $f(x, y) = x^2 + xy + y^2 + \frac{1}{x} + \frac{1}{y}$ , $x \neq 0, y \neq 0$ attains its local minimum value at the point $(a, b)$ , then the value of $a^3 + b^3$ is _____ (round off to TWO decimal places).

**GATE 2022 Mathematics (MA)**

Q.33	<p>If the ordinary differential equation</p> $x^2 \frac{d^2 \phi}{dx^2} + x \frac{d\phi}{dx} + x^2 \phi = 0, \quad x > 0$ <p>has a solution of the form <math>\phi(x) = x^r \sum_{n=0}^{\infty} a_n x^n</math>, where <math>a_n</math>'s are constants and <math>a_0 \neq 0</math>, then the value of <math>r^2 + 1</math> is _____.</p>

**GATE 2022 Mathematics (MA)**

Q.34	The Bessel functions $J_\alpha(x)$ , $x > 0$ , $\alpha \in \mathbb{R}$ satisfy $J_{\alpha-1}(x) + J_{\alpha+1}(x) = \frac{2\alpha}{x} J_\alpha(x)$ . Then, the value of $(\pi J_{\frac{3}{2}}(\pi))^2$ is _____.

Q.35	<p>The partial differential equation</p> $7\frac{\partial^2 u}{\partial x^2} + 16\frac{\partial^2 u}{\partial x \partial y} + 4\frac{\partial^2 u}{\partial y^2} = 0$ <p>is transformed to</p> $A\frac{\partial^2 u}{\partial \xi^2} + B\frac{\partial^2 u}{\partial \xi \partial \eta} + C\frac{\partial^2 u}{\partial \eta^2} = 0,$ <p>using <math>\xi = y - 2x</math> and <math>\eta = 7y - 2x</math>.</p> <p>Then, the value of <math>\frac{1}{12^3}(B^2 - 4AC)</math> is _____.</p>

**GATE 2022 Mathematics (MA)**

**Q.36 – Q.65 Carry TWO marks each.**

Q.36	Let $\mathbb{R}[X]$ denote the ring of polynomials in $X$ with real coefficients. Then, the quotient ring $\mathbb{R}[X]/(X^4 + 4)$ is
(A)	a field
(B)	an integral domain, but not a field
(C)	not an integral domain, but has 0 as the only nilpotent element
(D)	a ring which contains non-zero nilpotent elements

**GATE 2022 Mathematics (MA)**

Q.37	<p>Consider the following conditions on two proper non-zero ideals <math>J_1</math> and <math>J_2</math> of a non-zero commutative ring <math>R</math>.</p> <p><b>P:</b> For any <math>r_1, r_2 \in R</math>, there exists a unique <math>r \in R</math> such that <math>r - r_1 \in J_1</math> and <math>r - r_2 \in J_2</math>.</p> <p><b>Q:</b> <math>J_1 + J_2 = R</math></p> <p>Then, which of the following statements is TRUE?</p>
(A)	<b>P</b> implies <b>Q</b> but <b>Q</b> does not imply <b>P</b>
(B)	<b>Q</b> implies <b>P</b> but <b>P</b> does not imply <b>Q</b>
(C)	<b>P</b> implies <b>Q</b> and <b>Q</b> implies <b>P</b>
(D)	<b>P</b> does not imply <b>Q</b> and <b>Q</b> does not imply <b>P</b>

**GATE 2022 Mathematics (MA)**

Q.38	Let $f : [-\pi, \pi] \rightarrow \mathbb{R}$ be a continuous function such that $f(x) > \frac{f(0)}{2}$ , $ x  < \delta$ for some $\delta$ satisfying $0 < \delta < \pi$ . Define $P_{n,\delta}(x) = (1 + \cos x - \cos \delta)^n$ , for $n = 1, 2, 3, \dots$ . Then, which of the following statements is TRUE?
(A)	$\lim_{n \rightarrow \infty} \int_0^{2\delta} f(x) P_{n,\delta}(x) dx = 0$
(B)	$\lim_{n \rightarrow \infty} \int_{-2\delta}^0 f(x) P_{n,\delta}(x) dx = 0$
(C)	$\lim_{n \rightarrow \infty} \int_{-\delta}^{\delta} f(x) P_{n,\delta}(x) dx = 0$
(D)	$\lim_{n \rightarrow \infty} \int_{[-\pi, \pi] \setminus [-\delta, \delta]} f(x) P_{n,\delta}(x) dx = 0$

GATE 2022 Mathematics (MA)

Q.39	<p><b>P</b> : Suppose that <math>\sum_{n=0}^{\infty} a_n x^n</math> converges at <math>x = -3</math> and diverges at <math>x = 6</math>. Then <math>\sum_{n=0}^{\infty} (-1)^n a_n</math> converges.</p> <p><b>Q</b>: The interval of convergence of the series <math>\sum_{n=2}^{\infty} \frac{(-1)^n x^n}{4^n \log_e n}</math> is <math>[-4, 4]</math>.</p> <p>Which of the following statements is TRUE?</p>
(A)	<b>P</b> is true and <b>Q</b> is true
(B)	<b>P</b> is false and <b>Q</b> is false
(C)	<b>P</b> is true and <b>Q</b> is false
(D)	<b>P</b> is false and <b>Q</b> is true



**GATE 2022 Mathematics (MA)**

Q.40	<p>Let</p> $f_n(x) = \frac{x^2}{x^2 + (1 - nx)^2}, \quad x \in [0, 1], \quad n = 1, 2, 3, \dots$ <p>Then, which of the following statements is TRUE?</p>
(A)	$\{f_n\}$ is not equicontinuous on $[0, 1]$
(B)	$\{f_n\}$ is uniformly convergent on $[0, 1]$
(C)	$\{f_n\}$ is equicontinuous on $[0, 1]$
(D)	$\{f_n\}$ is uniformly bounded and has a subsequence converging uniformly on $[0, 1]$

**GATE 2022 Mathematics (MA)**

Q.41	Let $(\mathbb{Q}, d)$ be the metric space with $d(x, y) =  x - y $ . Let $E = \{p \in \mathbb{Q} : 2 < p^2 < 3\}$ . Then, the set $E$ is
(A)	closed but not compact
(B)	not closed but compact
(C)	compact
(D)	neither closed nor compact

**GATE 2022 Mathematics (MA)**

Q.42	Let $T : L^2[-1, 1] \rightarrow L^2[-1, 1]$ be defined by $Tf = \tilde{f}$ , where $\tilde{f}(x) = f(-x)$ almost everywhere. If $M$ is the kernel of $I - T$ , then the distance between the function $\phi(t) = e^t$ and $M$ is
(A)	$\frac{1}{2}\sqrt{(e^2 - e^{-2} + 4)}$
(B)	$\frac{1}{2}\sqrt{(e^2 - e^{-2} - 2)}$
(C)	$\frac{1}{2}\sqrt{(e^2 - 4)}$
(D)	$\frac{1}{2}\sqrt{(e^2 - e^{-2} - 4)}$

**GATE 2022 Mathematics (MA)**

Q.43	Let $X$ , $Y$ and $Z$ be Banach spaces. Suppose that $T : X \rightarrow Y$ is linear and $S : Y \rightarrow Z$ is linear, bounded and injective. In addition, if $S \circ T : X \rightarrow Z$ is bounded, then, which of the following statements is TRUE?
(A)	$T$ is surjective
(B)	$T$ is bounded but not continuous
(C)	$T$ is bounded
(D)	$T$ is not bounded

**GATE 2022 Mathematics (MA)**

Q.44	<p>The first derivative of a function <math>f \in C^\infty(-3, 3)</math> is approximated by an interpolating polynomial of degree 2, using the data</p> $(-1, f(-1)), (0, f(0)) \text{ and } (2, f(2)).$ <p>It is found that</p> $f'(0) \approx -\frac{2}{3}f(-1) + \alpha f(0) + \beta f(2).$ <p>Then, the value of <math>\frac{1}{\alpha\beta}</math> is</p>
(A)	3
(B)	6
(C)	9
(D)	12

**GATE 2022 Mathematics (MA)**

Q.45	The work done by the force $F = (x + y)\hat{i} - (x^2 + y^2)\hat{j}$ , where $\hat{i}$ and $\hat{j}$ are unit vectors in $\overrightarrow{OX}$ and $\overrightarrow{OY}$ directions, respectively, along the upper half of the circle $x^2 + y^2 = 1$ from $(1, 0)$ to $(-1, 0)$ in the $xy$ -plane is
(A)	$-\pi$
(B)	$-\frac{\pi}{2}$
(C)	$\frac{\pi}{2}$
(D)	$\pi$

**GATE 2022 Mathematics (MA)**

Q.46	<p>Let <math>u(x, t)</math> be the solution of the wave equation</p> $\frac{\partial^2 u}{\partial t^2} - \frac{\partial^2 u}{\partial x^2} = 0, \quad 0 < x < \pi, \quad t > 0,$ <p>with the initial conditions</p> $u(x, 0) = \sin x + \sin 2x + \sin 3x, \quad \frac{\partial u}{\partial t}(x, 0) = 0, \quad 0 < x < \pi$ <p>and the boundary conditions <math>u(0, t) = u(\pi, t) = 0, \quad t \geq 0</math>. Then, the value of <math>u\left(\frac{\pi}{2}, \pi\right)</math> is</p>
(A)	$-1/2$
(B)	$0$
(C)	$1/2$
(D)	$1$

**GATE 2022 Mathematics (MA)**

Q.47	<p>Let <math>T : \mathbb{R}^2 \rightarrow \mathbb{R}^2</math> be a linear transformation defined by</p> $T((1, 2)) = (1, 0) \quad \text{and} \quad T((2, 1)) = (1, 1).$ <p>For <math>p, q \in \mathbb{R}</math>, let <math>T^{-1}((p, q)) = (x, y)</math>.</p> <p>Which of the following statements is TRUE?</p>
(A)	$x = p - q; y = 2p - q$
(B)	$x = p + q; y = 2p - q$
(C)	$x = p + q; y = 2p + q$
(D)	$x = p - q; y = 2p + q$



# GATE 2022 Mathematics (MA)

Q.48	<p>Let <math>y = (\alpha, -1)^T</math>, <math>\alpha \in \mathbb{R}</math> be a feasible solution for the dual problem of the linear programming problem</p> $\begin{aligned} &\text{Maximize:} && 5x_1 + 12x_2 \\ &\text{subject to:} && x_1 + 2x_2 + x_3 \leq 10 \\ &&& 2x_1 - x_2 + 3x_3 = 8 \\ &&& x_1, x_2, x_3 \geq 0. \end{aligned}$ <p>Which of the following statements is TRUE?</p>
(A)	$\alpha < 3$
(B)	$3 \leq \alpha < 5.5$
(C)	$5.5 \leq \alpha < 7$
(D)	$\alpha \geq 7$

**GATE 2022 Mathematics (MA)**

Q.49	Let $K$ denote the subset of $\mathbb{C}$ consisting of elements algebraic over $\mathbb{Q}$ . Then, which of the following statements are TRUE?
(A)	No element of $\mathbb{C} \setminus K$ is algebraic over $\mathbb{Q}$
(B)	$K$ is an algebraically closed field
(C)	For any bijective ring homomorphism $f : \mathbb{C} \rightarrow \mathbb{C}$ , we have $f(K) = K$
(D)	There is no bijection between $K$ and $\mathbb{Q}$

**GATE 2022 Mathematics (MA)**

Q.50	Let $T$ be a Möbius transformation such that $T(0) = \alpha$ , $T(\alpha) = 0$ and $T(\infty) = -\alpha$ , where $\alpha = (-1 + i)/\sqrt{2}$ . Let $L$ denote the straight line passing through the origin with slope $-1$ , and let $C$ denote the circle of unit radius centred at the origin. Then, which of the following statements are TRUE?
(A)	$T$ maps $L$ to a straight line
(B)	$T$ maps $L$ to a circle
(C)	$T^{-1}$ maps $C$ to a straight line
(D)	$T^{-1}$ maps $C$ to a circle

**GATE 2022 Mathematics (MA)**

Q.51	Let $a > 0$ . Define $D_a : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$ by $(D_a f)(x) = \frac{1}{\sqrt{a}} f\left(\frac{x}{a}\right)$ , almost everywhere, for $f \in L^2(\mathbb{R})$ . Then, which of the following statements are TRUE?
(A)	$D_a$ is a linear isometry
(B)	$D_a$ is a bijection
(C)	$D_a \circ D_b = D_{a+b}$ , $b > 0$
(D)	$D_a$ is bounded from below

**GATE 2022 Mathematics (MA)**

Q.52	Let $\{\phi_0, \phi_1, \phi_2, \dots\}$ be an orthonormal set in $L^2[-1, 1]$ such that $\phi_n = C_n P_n$ , where $C_n$ is a constant and $P_n$ is the Legendre polynomial of degree $n$ , for each $n \in \mathbb{N} \cup \{0\}$ . Then, which of the following statements are TRUE?
(A)	$\phi_6(1) = 1$
(B)	$\phi_7(-1) = 1$
(C)	$\phi_7(1) = \sqrt{\frac{15}{2}}$
(D)	$\phi_6(-1) = \sqrt{\frac{13}{2}}$

**GATE 2022 Mathematics (MA)**

Q.53	Let $X = (\mathbb{R}, T)$ , where $T$ is the smallest topology on $\mathbb{R}$ in which all the singleton sets are closed. Then, which of the following statements are TRUE?
(A)	$[0, 1]$ is compact in $X$
(B)	$X$ is not first countable
(C)	$X$ is second countable
(D)	$X$ is first countable

**GATE 2022 Mathematics (MA)**

Q.54	<p>Consider <math>(\mathbb{Z}, T)</math>, where <math>T</math> is the topology generated by sets of the form</p> $A_{m,n} = \{m + nk \mid k \in \mathbb{Z}\},$ <p>for <math>m, n \in \mathbb{Z}</math> and <math>n \neq 0</math>. Then, which of the following statements are TRUE?</p>
(A)	$(\mathbb{Z}, T)$ is connected
(B)	Each $A_{m,n}$ is a closed subset of $(\mathbb{Z}, T)$
(C)	$(\mathbb{Z}, T)$ is Hausdorff
(D)	$(\mathbb{Z}, T)$ is metrizable

**GATE 2022 Mathematics (MA)**

Q.55	<p>Let <math>A \in \mathbb{R}^{m \times n}</math>, <math>c \in \mathbb{R}^n</math> and <math>b \in \mathbb{R}^m</math>. Consider the linear programming primal problem</p> $\begin{aligned} &\text{Minimize: } c^T x \\ &\text{subject to: } Ax = b \\ &\quad \quad \quad x \geq 0. \end{aligned}$ <p>Let <math>x^0</math> and <math>y^0</math> be feasible solutions of the primal and its dual, respectively. Which of the following statements are TRUE?</p>
(A)	$c^T x^0 \geq b^T y^0$
(B)	$c^T x^0 = b^T y^0$
(C)	If $c^T x^0 = b^T y^0$ , then $x^0$ is optimal for the primal
(D)	If $c^T x^0 = b^T y^0$ , then $y^0$ is optimal for the dual



**GATE 2022 Mathematics (MA)**

Q.56	<p>Consider <math>\mathbb{R}^3</math> as a vector space with the usual operations of vector addition and scalar multiplication. Let <math>x \in \mathbb{R}^3</math> be denoted by <math>x = (x_1, x_2, x_3)</math>. Define subspaces <math>W_1</math> and <math>W_2</math> by</p> $W_1 := \{x \in \mathbb{R}^3 : x_1 + 2x_2 - x_3 = 0\}$ <p>and</p> $W_2 := \{x \in \mathbb{R}^3 : 2x_1 + 3x_3 = 0\}.$ <p>Let <math>\dim(U)</math> denote the dimension of the subspace <math>U</math>.</p> <p>Which of the following statements are TRUE?</p>
(A)	$\dim(W_1) = \dim(W_2)$
(B)	$\dim(W_1) + \dim(W_2) - \dim(\mathbb{R}^3) = 1$
(C)	$\dim(W_1 + W_2) = 2$
(D)	$\dim(W_1 \cap W_2) = 1$

**GATE 2022 Mathematics (MA)**

Q.57	<p>Three companies <math>C_1, C_2</math> and <math>C_3</math> submit bids for three jobs <math>J_1, J_2</math> and <math>J_3</math>. The costs involved per unit are given in the table below:</p> <table><tr><td></td><td><math>J_1</math></td><td><math>J_2</math></td><td><math>J_3</math></td></tr><tr><td><math>C_1</math></td><td>10</td><td>12</td><td>8</td></tr><tr><td><math>C_2</math></td><td>9</td><td>15</td><td>10</td></tr><tr><td><math>C_3</math></td><td>15</td><td>10</td><td>9</td></tr></table> <p>Then, the cost of the optimal assignment is _____.</p>		$J_1$	$J_2$	$J_3$	$C_1$	10	12	8	$C_2$	9	15	10	$C_3$	15	10	9
	$J_1$	$J_2$	$J_3$														
$C_1$	10	12	8														
$C_2$	9	15	10														
$C_3$	15	10	9														

**GATE 2022 Mathematics (MA)**

Q.58	<p>The initial value problem <math>\frac{dy}{dx} = f(x, y)</math>, <math>y(x_0) = y_0</math> is solved by using the following second order Runge-Kutta method:</p> $\begin{aligned} K_1 &= hf(x_i, y_i) \\ K_2 &= hf(x_i + \alpha h, y_i + \beta K_1) \\ y_{i+1} &= y_i + \frac{1}{4}(K_1 + 3K_2), \quad i \geq 0, \end{aligned}$ <p>where <math>h</math> is the uniform step length between the points <math>x_0, x_1, \dots, x_n</math> and <math>y_i = y(x_i)</math>. The value of the product <math>\alpha\beta</math> is _____ (round off to TWO decimal places).</p>

**GATE 2022 Mathematics (MA)**

Q.59	The surface area of the paraboloid $z = x^2 + y^2$ between the planes $z = 0$ and $z = 1$ is _____ (round off to ONE decimal place).

**GATE 2022 Mathematics (MA)**

Q.60	The rate of change of $f(x, y, z) = x + x \cos z - y \sin z + y$ at $P_0$ in the direction from $P_0(2, -1, 0)$ to $P_1(0, 1, 2)$ is _____.

**GATE 2022 Mathematics (MA)**

Q.61	<p>If the Laplace equation</p> $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0, \quad 1 < x < 2, \quad 1 < y < 2$ <p>with the boundary conditions</p> $\frac{\partial u}{\partial x}(1, y) = y, \quad \frac{\partial u}{\partial x}(2, y) = 5, \quad 1 < y < 2$ <p>and</p> $\frac{\partial u}{\partial y}(x, 1) = \frac{\alpha x^2}{7}, \quad \frac{\partial u}{\partial y}(x, 2) = x, \quad 1 < x < 2$ <p>has a solution, then the constant <math>\alpha</math> is _____.</p>

**GATE 2022 Mathematics (MA)**

Q.62	<p>Let <math>u(x, y)</math> be the solution of the first order partial differential equation</p> $x \frac{\partial u}{\partial x} + (x^2 + y) \frac{\partial u}{\partial y} = u, \text{ for all } x, y \in \mathbb{R}$ <p>satisfying <math>u(2, y) = y - 4, y \in \mathbb{R}</math>. Then, the value of <math>u(1, 2)</math> is _____.</p>

**GATE 2022 Mathematics (MA)**

Q.63	<p>The optimal value for the linear programming problem</p> $\begin{aligned} \text{Maximize: } & 6x_1 + 5x_2 \\ \text{subject to: } & 3x_1 + 2x_2 \leq 12 \\ & -x_1 + x_2 \leq 1 \\ & x_1, x_2 \geq 0 \end{aligned}$ <p>is _____.</p>



# GATE 2022 Mathematics (MA)

Q.64	<p>A certain product is manufactured by plants <math>P_1, P_2</math> and <math>P_3</math> whose capacities are 15, 25 and 10 units, respectively. The product is shipped to markets <math>M_1, M_2, M_3</math> and <math>M_4</math>, whose requirements are 10, 10, 10 and 20, respectively. The transportation costs per unit are given in the table below.</p> <table><tr><td></td><td><math>M_1</math></td><td><math>M_2</math></td><td><math>M_3</math></td><td><math>M_4</math></td><td></td></tr><tr><td><math>P_1</math></td><td>1</td><td>3</td><td>1</td><td>3</td><td>15</td></tr><tr><td><math>P_2</math></td><td>2</td><td>2</td><td>4</td><td>1</td><td>25</td></tr><tr><td><math>P_3</math></td><td>2</td><td>1</td><td>1</td><td>2</td><td>10</td></tr><tr><td></td><td>10</td><td>10</td><td>10</td><td>20</td><td></td></tr></table> <p>Then the cost corresponding to the starting basic solution by the Northwest-corner method is _____.</p>		$M_1$	$M_2$	$M_3$	$M_4$		$P_1$	1	3	1	3	15	$P_2$	2	2	4	1	25	$P_3$	2	1	1	2	10		10	10	10	20	
	$M_1$	$M_2$	$M_3$	$M_4$																											
$P_1$	1	3	1	3	15																										
$P_2$	2	2	4	1	25																										
$P_3$	2	1	1	2	10																										
	10	10	10	20																											

**GATE 2022 Mathematics (MA)**

Q.65	Let $M$ be a $3 \times 3$ real matrix such that $M^2 = 2M + 3I$ . If the determinant of $M$ is $-9$ , then the trace of $M$ equals _____.

Q. No.	Session	Question Type	Subject Name	Key/Range	Mark
1	2	MCQ	GA	D	1
2	2	MCQ	GA	A	1
3	2	MCQ	GA	A	1
4	2	MCQ	GA	A	1
5	2	MCQ	GA	C	1
6	2	MCQ	GA	C	2
7	2	MCQ	GA	A	2
8	2	MCQ	GA	B	2
9	2	MCQ	GA	A	2
10	2	MCQ	GA	C	2
11	2	MCQ	MA	D	1
12	2	MCQ	MA	C	1
13	2	MCQ	MA	C	1
14	2	MCQ	MA	D	1
15	2	MCQ	MA	A	1
16	2	MCQ	MA	D	1
17	2	MCQ	MA	C	1
18	2	MCQ	MA	B	1
19	2	MCQ	MA	A	1
20	2	MCQ	MA	A	1
21	2	MCQ	MA	B	1
22	2	MCQ	MA	B	1
23	2	MCQ	MA	D	1
24	2	MCQ	MA	B	1
25	2	MSQ	MA	B, C, D	1
26	2	NAT	MA	30 to 30	1
27	2	NAT	MA	6 to 6	1
28	2	NAT	MA	0.55 to 0.59	1
29	2	NAT	MA	5 to 5	1
30	2	NAT	MA	1 to 1	1
31	2	NAT	MA	39 to 39	1
32	2	NAT	MA	0.65 to 0.68	1
33	2	NAT	MA	1 to 1	1
34	2	NAT	MA	2 to 2	1
35	2	NAT	MA	12 to 12	1
36	2	MCQ	MA	C	2
37	2	MCQ	MA	A	2
38	2	MCQ	MA	D	2
39	2	MCQ	MA	C	2
40	2	MCQ	MA	A	2
41	2	MCQ	MA	A	2
42	2	MCQ	MA	D	2
43	2	MCQ	MA	C	2
44	2	MCQ	MA	D	2

45	2	MCQ	MA	B	2
46	2	MCQ	MA	B	2
47	2	MCQ	MA	B	2
48	2	MCQ	MA	D	2
49	2	MSQ	MA	A, B, C	2
50	2	MSQ	MA	A, C	2
51	2	MSQ	MA	A, B, D	2
52	2	MSQ	MA	C, D	2
53	2	MSQ	MA	A, B	2
54	2	MSQ	MA	B, C, D	2
55	2	MSQ	MA	A, C, D	2
56	2	MSQ	MA	A, B, D	2
57	2	NAT	MA	27 to 27	2
58	2	NAT	MA	0.43 to 0.45	2
59	2	NAT	MA	5.1 to 5.5	2
60	2	NAT	MA	0 to 0	2
61	2	NAT	MA	15 to 15	2
62	2	NAT	MA	1 to 1	2
63	2	NAT	MA	27 to 27	2
64	2	NAT	MA	105 to 105	2
65	2	NAT	MA	5 to 5	2